

CHAPTRE TWO


VECTORS

- A vector is a physical quantity which has both magnitude and direction.
- Example are
 - a. A force of 20N acting North.
 - b. A velocity of 5km/h East.

Types of vectors:


- In general there are two types and these are
 - i. Free vector.
 - ii. Position vector.

Free vector:



- A free vector is a vector which does not pass through any specific position.
- They are usually represented by small letters e.g. \vec{e} , \vec{g} , \vec{a} , \vec{b} and \vec{c}


Position vector :



This is a vector which passes through the origin or a specified point.


Vector notation:

- A vector may be represented by a line segment as shown next:



- This given vector can be represented by \overrightarrow{AB} , \overline{AB} , \underline{AB} , \widetilde{AB} , AB .

The Triangle law:



According to the triangle law, $\overrightarrow{AC} = \overrightarrow{AB} + \overrightarrow{BC} \Rightarrow \overrightarrow{AB} = \overrightarrow{AC} - \overrightarrow{BC}$ and $\overrightarrow{BC} = \overrightarrow{AC} - \overrightarrow{AB}$

The unit vector:

- This is a vector whose magnitude is one in the direction under consideration.
- The unit vector along a vector \vec{a} is written as \hat{a}
- Also the unit vector along a vector \vec{b} is written as \hat{b}
- The unit vector along the vector \overrightarrow{BC} is written as \widehat{BC}
- Consider the vector $\overrightarrow{A \rightarrow B} = 1$
- The vector is written as \overrightarrow{AB} and its unit vector is written as \widehat{AB} .

Equal vectors:

- Two vectors are said to be equal if their magnitudes and directions are equal
- Example are $\overrightarrow{AB} = 50\text{km/h E}$ and $\overrightarrow{CD} = 50\text{km/h E}$.

The negative vector:


- The negative of the vector \vec{a} is written as $-\vec{a}$
- If $-\vec{a}$ is the negative vector of the vector \vec{a} , then $\vec{a} + (-\vec{a}) = \vec{0}$.
- The vector $-\vec{a}$ is a vector of the same magnitude as \vec{a} , but it is opposite in direction.
- It must be noted that $\overrightarrow{AB} + \overrightarrow{BA} = \vec{0}$.
- Also if $\vec{b} = \overrightarrow{CD}$, then $-\vec{b} = \overrightarrow{DC}$, and $\overrightarrow{CD} + \overrightarrow{DC} = \vec{0}$.
- If we consider a vector \overrightarrow{CD} , then its negative vector is \overrightarrow{DC} .

The zero vector (null vector):

- This is a vector where magnitude is zero and its direction is undefined.
- It is represented by $\vec{0} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$

Notation of the magnitude of a vectors:


- If \overrightarrow{AB} is a vector, then its magnitude is written as $|\overrightarrow{AB}|$
- Similarly the magnitude of the vector \vec{b} is written as $|\vec{b}|$
- If $\overrightarrow{OP} = \begin{pmatrix} a \\ b \end{pmatrix}$, then its magnitude $= |\overrightarrow{OP}| = \sqrt{a^2 + b^2}$



Q1. i. If $OP = \begin{pmatrix} 6 \\ 5 \end{pmatrix}$, find the magnitude of \overrightarrow{OP} .

ii. Find ϕ the angle between \overrightarrow{OP} and the x-axis

Soln.



i. $|\overrightarrow{OP}| = \sqrt{6^2 + 5^2} = \sqrt{61} = 7.8$

ii. $\tan \phi = \frac{5}{6} \Rightarrow \tan \phi = 0.83 \Rightarrow \phi = \tan^{-1} 0.83 \Rightarrow \phi = 40.$


Scalar multiplication of vector:

- If λ is the scalar and \vec{a} is the vector, then the scalar \times the vector $= \lambda \vec{a}$
- When a scalar multiplies a vector, the product is also a vector, and for this reason $\lambda \vec{a}$ is also a vector.
- The vector $\lambda \vec{a}$ is parallel to \vec{a} , and is in the same direction as \vec{a} , but has λ times the magnitude of \vec{a} .
- For example the vectors \vec{a} and $2\vec{a}$ have the same direction.

i.e. \vec{a} $\quad \quad \quad 2\vec{a}$

- But the vectors \vec{a} and $-2\vec{a}$ are opposite in direction.
- $\lambda(\vec{a} + \vec{b}) = \lambda \vec{a} + \lambda \vec{b}$, e.g. $6(\vec{a} + \vec{b}) = 6\vec{a} + 6\vec{b}$
- Also $(2 + 4)\vec{a} = 2\vec{a} + 4\vec{a}$
- Finally $\lambda_1(\lambda_2 \vec{a}) = \lambda_1 \lambda_2 \vec{a}$, e.g. $3(2\vec{a}) = 6\vec{a}$

N/B: y



$$\leftarrow O \rightarrow x_1$$

- If $P(x_1, y_1)$ is a point in the $x - y$ plane, then the position vector of P relative to the origin, O is defined by $\overrightarrow{OP} = \begin{pmatrix} x_1 \\ y_1 \end{pmatrix}$
- Also if $A = (0, 6)$, then $\overrightarrow{OA} = \begin{pmatrix} 0 \\ 6 \end{pmatrix}$

Q2. Find the numbers m and n such that

$$M \begin{pmatrix} 3 \\ 5 \end{pmatrix} + n \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 4 \\ 9 \end{pmatrix}$$

Soln.

$$M \begin{pmatrix} 3 \\ 5 \end{pmatrix} + n \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 4 \\ 9 \end{pmatrix} \Rightarrow \begin{pmatrix} 3m \\ 5m \end{pmatrix} + \begin{pmatrix} 2n \\ n \end{pmatrix} = \begin{pmatrix} 4 \\ 9 \end{pmatrix}$$

$$\Rightarrow 3m + 2n = 4 \dots \dots \text{eqn(1).}$$

$$5m + n = 9 \dots \dots \dots \text{eqn(2)}$$

Solve eqns (1) and (2) simultaneously

$$\Rightarrow m = 2 \text{ and } n = -1$$

Q3. If $mp + nq = \begin{pmatrix} 4 \\ 3 \end{pmatrix}$, find m and n where m and n are scalar, given that $p = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$ and $q = \begin{pmatrix} 2 \\ 5 \end{pmatrix}$

Soln.

$$p = \begin{pmatrix} 2 \\ 3 \end{pmatrix} \text{ and } q = \begin{pmatrix} 2 \\ 5 \end{pmatrix} \text{ but } mp + nq = \begin{pmatrix} 4 \\ 3 \end{pmatrix}$$

$$\Rightarrow m \begin{pmatrix} 2 \\ 3 \end{pmatrix} + n \begin{pmatrix} 2 \\ 5 \end{pmatrix} = \begin{pmatrix} 4 \\ 3 \end{pmatrix} \Rightarrow \begin{pmatrix} 2m \\ 3m \end{pmatrix} + \begin{pmatrix} 2n \\ 5n \end{pmatrix} = \begin{pmatrix} 4 \\ 3 \end{pmatrix}$$

$$\Rightarrow 2m + 2n = 4 \dots (1)$$

$$3m + 5n = 3 \dots (2)$$

Solve eqns (1) and (2) simultaneously to get the values of m and n .

Q4. If $r = \begin{pmatrix} 3 \\ 1 \end{pmatrix}$ and $s = \begin{pmatrix} -2 \\ 1 \end{pmatrix}$, evaluate $6(r + 2s)$

Soln.

$$\text{Consider } 6(r + 2s), \text{ solve what is inside the bracket first } \Rightarrow r + 2s = \begin{pmatrix} 3 \\ 1 \end{pmatrix} + \begin{pmatrix} -2 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 1 \end{pmatrix} + 2 \begin{pmatrix} -2 \\ 1 \end{pmatrix} \Rightarrow r + 2s = \begin{pmatrix} 3 + (-4) \\ 1 + 2 \end{pmatrix} = \begin{pmatrix} -1 \\ 3 \end{pmatrix} \Rightarrow 6(r + 2s) = 6 \begin{pmatrix} -1 \\ 3 \end{pmatrix} = \begin{pmatrix} -6 \\ 18 \end{pmatrix}$$

Q5. If $p = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$, $q = \begin{pmatrix} -2 \\ 3 \end{pmatrix}$ and $r = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$, find $2p - q + r$

Soln.

$$2p - q + r = 2 \begin{pmatrix} 1 \\ 2 \end{pmatrix} - \begin{pmatrix} -2 \\ 3 \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 4 \end{pmatrix} - \begin{pmatrix} -2 \\ 3 \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 + 2 + 1 \\ 4 - 3 + 1 \end{pmatrix} = \begin{pmatrix} 5 \\ 2 \end{pmatrix} \Rightarrow 2p - q + r = \begin{pmatrix} 5 \\ 2 \end{pmatrix}.$$